

10018604

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A23L 1/212, 1/06, 1/23, 3/015</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/38591</b> <b>(43) International Publication Date:</b> 23 October 1997 (23.10.97)
<b>(21) International Application Number:</b> PCT/EP97/01265 <b>(22) International Filing Date:</b> 11 March 1997 (11.03.97) <b>(30) Priority Data:</b> 96302608.3 12 April 1996 (12.04.96) EP <i>(34) Countries for which the regional or international application was filed:</i> GB et al. <b>(71) Applicant (for AU BB CA GB IE IL KE LC LK LS MN MW NZ SD SG SZ TT UG only):</b> UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). <b>(71) Applicant (for all designated States except AU BB CA GB IE IL KE LC LK LS MN MW NZ SD SG SZ TT UG):</b> UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). <b>(72) Inventors:</b> WILDING, Peter; 29 Dryden Street, Raunds, Wellingborough, Northamptonshire NN9 5EL (GB). WOOLNER, Elizabeth, Mary; 41 Whitley Road, Bedford MK42 0XA (GB). <b>(74) Agent:</b> EKE, Philippa, Dianne; Unilever plc, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>

**(54) Title:** PROCESS FOR THE PREPARATION OF A FOOD PRODUCT**(57) Abstract**

A process for providing a tomato-based product comprising the following steps: a) applying ultra high pressure to a tomato piece, such that polygalacturonase is inactivated and pectinmethylesterase and preferably peroxidase are not inactivated; b) incubating the tomato piece with endogenous pectinmethylesterase to achieve a desired consistency and preferably with peroxidase to achieve a fresh aroma and/or flavour profile; and c) inactivating the pectinmethylesterase and any peroxidase.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR THE PREPARATION OF A FOOD  
PRODUCT

Field of the Invention

5

The invention relates to a process for the preparation of tomato-based products of high quality and consistency and products obtained by this process.

10

Background to the Invention

15

It is desirable to be able to produce tomato-based products having a high consistency and quality with a low solids content. In order to do this, it is necessary to be able to substantially increase the viscosity of the tomato pastes commonly used to prepare products such as sauces.

20

It is also desirable to be able to produce tomato-based products having a fresh, rather than a processed, flavour and/or aroma profile.

25

Enzymes, such as polygalacturonase (PG) and pectinmethylesterase (PME), are naturally-present in tomatoes. If the tomatoes are broken-down, these enzymes are released and pectinmethylesterase catalyses the removal of methyl ester groups from pectin molecules occurring naturally in the tomatoes, thereby producing pectic and pectinic acids; polygalacturonase rapidly depolymerises these pectic and pectinic acids. Consequently, the tomato-based product lacks pectic substances and has a thin and watery consistency.

30

35

Heating a paste of broken-down tomatoes above 80°C inactivates PME and PG, thereby preventing breakdown of pectic substances in tomato products during processing, such that a good consistency is maintained. However, heating leads to increased rates of chemical reactions

including the Maillard reaction, which leads to the browning of tomato products and the generation of off-flavours.

5 High Pressure and Biotechnology, Colloque INSERM/1992, Vol 24 pp 195-209 discusses the effects of high pressure (1-7 kbar) on enzymes. It states that polyphenoloxidase does not appear to be inactivated by UHP. It also mentions the conditions necessary to inactivate PME in citrus fruits, and concludes that pressurisation conditions sufficient for the microbial 'pasteurisation' of fruit juices or purees do not fully stabilise them against modifications by PME and some other enzymes. These 'pressure-pasteurised' acid foods therefore also require a mild thermal blanching, a refrigerated storage or the addition of enzyme inhibitors, to fully inactivate the degrading enzymes. Enzyme inactivation is important for achieving long term ambient stability.

20 The present invention seeks to provide a process for manipulating the consistency and texture of tomato-based products having a low solids content. Preferably, the tomato-based product has a flavour/aroma profile and a colour which is perceived as fresher than conventionally processed tomato-based products.

#### Summary of the Invention

30 According to the present invention there is provided a process for providing a tomato-based product, comprising the following steps:

- 35 a) applying ultra high pressure to a tomato piece, such that polygalacturonase is inactivated and pectinmethylesterase is not inactivated;
- b) incubating the tomato piece with endogenous

pectinmethylesterase to achieve a desired consistency;  
and

c) inactivating the pectinmethylesterase.

5

The inventors of the present invention were surprised to find that under suitable conditions, polygalacturonase is inactivated by applying ultra-high pressure (UHP), whereas pectinmethylesterase remains active. For example, applying  
10 a pressure of 500MPa for 15 minutes at approximately 20°C achieves this selective inactivation of PG.

In the absence of polygalacturonase to depolymerise the pectic and pectinic acids formed by the action of PME on pectin, these acids remain in the cell wall to prevent  
15 structural breakdown. The selective inactivation of polygalacturonase may therefore prevent texture degradation.

Also, incubating the tomato piece with endogenous PME in step b results in a firming effect and an improvement in texture. This is because the removal of methyl ester groups from pectin molecules, catalysed by PME, results in the pectin molecules being able to associate with each  
20 other via cation cross-links to provide an increase in consistency. The incubation may be conducted at 42°C for 30 minutes.

Once the desired thickness has been achieved, the PME is  
30 inactivated by, for example, heating, adjusting pH and/or adding inhibitors. If heating is used, the temperature is typically raised to 90°C for 10 minutes. Such a heat treatment is sufficient to pasteurise the product. Hence, a single heating step may be used both to inactivate PME  
35 and to pasteurise, resulting in a treatment which requires less heat input in comparison to the use of separate inactivation and pasteurisation heating steps.

The inventors have also found that, under certain UHP conditions, the enzyme peroxidase remains active together with PME after PG has been inactivated. They have found that peroxidase is more stable to UHP than PG, but less stable than PME. In contrast, the inventors have found that peroxidase is less heat stable than both PG and PME; therefore, peroxidase is inactivated before PG and PME in the heat treatments of the processes conventionally used for preparing tomato products.

Without wishing to be bound by theory, the inventors believe that peroxidase is important in the generation of volatiles for providing a fresh tomato flavour/aroma profile, and that the presence of active peroxidase in the tomato product during the incubation step b is advantageous to the perception of freshness in the final tomato product. The process of the present invention may therefore result in a tomato product having a flavour/aroma profile which is perceived as fresher than conventionally processed tomato products, since peroxidase is inactivated relatively quickly during the heating of conventional processes.

In the present invention, any active peroxidase is inactivated in step c, together with PME, thus preventing the flavour and/or aroma effects developing to extremes and decreasing quality.

A tomato piece is, for example, whole tomatoes either peeled or unpeeled, or a piece of chopped, diced, comminuted or macerated tomato. The tomato piece may have a size in the range of from approximately 10  $\mu$ m to approximately 10cm and may form part of a tomato paste, puree or juice. Commonly, the present invention is applied to a plurality of tomato pieces simultaneously.

An advantage of the process of the present invention is the manipulation of the viscosity and texture of tomato-based

products. Once PG has been inactivated using UHP, viscosity and texture is controlled by the conditions of the incubation step b and the inactivation step c; ie. they control the degree of firming which occurs. Moreover, using the process of the present invention, the tomato-based products preferably have a flavour/aroma profile and/or a colour which is perceived as fresher than conventionally processed tomato-based products. Therefore, by selecting the right conditions for steps a, b and c, viscosity and/or texture improvement, and preferably fresh colour, flavour and/or aroma generation, can be achieved.

Thus, the present invention provides means for selectively inactivating deleterious enzymes, and for maintaining and promoting the activity of other enzymes to control and improve viscosity and texture, and, preferably, also to improve colour, flavour and/or aroma.

Examples of the products and processes of the invention will now be described to illustrate, but not to limit, the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

##### Example 1

Eight 150g samples of tomato dice were subjected to a range of ultra high pressures for different lengths of time using a high hydrostatic pressure unit. The pressure and time treatments used are listed in Table 1.

Pressure, up to a maximum of 600MPa, was provided by water pumped into the base of the unit. Water was used as the pressure transferring medium. Generally, it took 2-3 minutes for the UHP system to reach the chosen pressure. It also took 2-3 mins to depressurize after treatment. The pressure and temperature, measured at four points within

the unit, were logged continuously by a computer. All experiments were carried out at ambient temperature.

5 A portion of each pressure-treated sample was stored for analysis either by freezing it rapidly using dry ice and storing at  $-20^{\circ}\text{C}$  or by packing it in 150g of tomato juice in foil pouches, heat sealing the pouches and heat treating for 30 minutes in a waterbath set at  $97^{\circ}\text{C}$ . This heat treatment was sufficient to inactivate all enzymes and to  
10 achieve pasteurisation.

The remainder of each pressure-treated sample was placed in a sachet and incubated for 30 minutes in a waterbath set at either  $40^{\circ}\text{C}$  or  $60^{\circ}\text{C}$ .

15 These samples were also stored for analysis, using the method described above.

20 For each stored sample, the activity of the enzymes pectin methyl esterase (PME), polygalacturonase (PG), lipoxygenase and peroxidase was determined using methods i to iv described below. The activity of these enzymes was also measured in a sample of the tomato dice which had not undergone the pressure and thermal incubation treatments  
25 described above (this was the control sample).

i) PME activity was assayed by adding  $200\mu\text{l}$  of sample to 20mls 1% apple pectin in 0.15M sodium chloride (pH 7.5 with sodium hydroxide). Activity was calculated from the slope  
30 of mls 0.025M NaOH used to maintain pH 7.5 at  $30^{\circ}\text{C}$  against time.

35 ii) PG activity was measured by the detection of reducing ends, which are generated when the enzyme breaks down polygalacturonic acid, using the colorimetric PAHBAH assay (Lever M. (1972) Anal Biochem 47: 273-279). PAHBAH is an abbreviation of the chemical p-hydroxybenzoic acid



hydrazide, which is used for the colour detection of reducing sugars. A high salt extract was prepared by adding low salt buffer (25mM sodium acetate buffer pH 5.6) to a tomato sample at a ratio of 1:2 volume:weight, shaking vigorously then centrifuging at 10,000g for 20 minutes at 4°C. After centrifugation, the supernatant was poured away and the pellet resuspended in an equivalent volume of a high salt buffer (25mM sodium acetate buffer pH 5.6 containing 1.0M sodium chloride) and left to stand at 4°C for 1 hour. The suspension was then respun at 10,000g, 4°C for 20 minutes and the supernatant (high salt extract) stored at -70°C in 1ml aliquots.

For PG assays, the high salt extract was diluted 1:100 in 50mM sodium acetate buffer pH 4.0 containing 0.2M NaCl; the activity was assayed using 0.2% polygalacturonic acid in a total assay volume of 500 $\mu$ l. The assays were incubated at 40°C for 1 hour and terminated by generating colour, which comprised adding 5mls of PAHBAH reagent and boiling for 6 minutes. After cooling under running water to room temperature, absorbance was read at 410nm using quartz cuvettes against a standard curve of galacturonic acid. Controls were performed by terminating assays with PAHBAH reagent as soon as the enzyme extract was added, ie. at zero time, and boiling as above.

iii) Lipxygenase activity was determined at 30°C using the polarographic method of Nicolas et al (1982 J Sci Food Agric 33:365-372) using linoleic acid (28mM final concentration) dispersed in 0.1M sodium phosphate buffer pH 7.0 containing polyoxyethylenesorbitan monolaurate (0.8% final concentration) in a total volume of 5ml.

iv) Peroxidase activity was determined by the spectrophotometric quantification of o-quinone production in a mixture containing 3.0 $\mu$ l 30% hydrogen peroxide (8.8mM final concentration), phosphate buffer pH 7.0 and enzyme

extract, in a total volume of 2.925ml. The reaction was initiated by adding 0.075ml of guaiacol (5mM final concentration) and measuring the change in absorbance at 400nm using a spectrophotometer with a temperature controlled cuvette holder at 30°C.

The results of these measurements for each sample after pressure-treatment alone are shown in Table 1 below.

Table 1

Treatment	% activity remaining after treatment			
	PME	PG	Lipoxygenase	Peroxidase
300MPa, 5 min	100	100	-	-
300MPa, 15 min	100	99	-	-
400MPa, 10 min	100	100	74	96
400MPa, 20 min	100	62	51	79
500MPa, 10 min	100	3	1	77
500MPa, 20 min	100	0	0	71
600MPa, 10 min	100	0	0	5
600MPa, 20 min	100	0	0	0

The results of these measurements for samples after pressure-treatment alone and after both pressure-treatment and thermal incubation are shown in Table 2 below, together with the results for the control sample.

Table 2

Treatment	% activity remaining after treatment			
	PME	PG	Lipoxye nase	Peroxid ase
No treatment (control)	100	100	100	100
400MPa 10 mins	100	100	74	96
400MPa 10 mins + 40°C 30 mins	90	76	33	100
400MPa 10 mins + 60°C 30 mins	58	45	0	100
600MPa 20 mins	100	0	0	0
600MPa 20 mins + 40°C 30 mins	86	0	0	0
600MPa 20 mins + 60°C 30 mins	72	0	0	0

The results in Table 1 show that PME is the most barostable enzyme. Its activity is not affected after a pressure treatment of 600MPa for 20 minutes. Peroxidase is the next most stable enzyme, followed by PG and lipoxxygenase, which have similar stabilities. After a pressure treatment of 500MPa for 20 mins, both PME and peroxidase are active whilst PG and lipoxxygenase have been inactivated; the presence of peroxidase is favourable to flavour and aroma.

From Table 2, it is clear that there are pressures which inactivate PG, lipoxxygenase and peroxidase but do not inactivate PME. Subsequent incubation allows the active

PME to promote firming in the absence of PG-promoted texture degradation, as evidenced by example 2 below.

Example 2

5

10 In accordance with the methods of example 1, tomato juice samples were treated with UHP (either 400MPa for 10 mins which is not sufficient for PG inactivation or 600MPa for 20 mins which is sufficient for PG inactivation) and incubated at either 40 or 60°C for 30 mins. Samples were then pasteurized by filling into foil pouches, heat sealing and placing in a waterbath at 95°C for 30 minutes. All enzyme activities were inactivated after this pasteurising heat treatment.

15

20 The effects of the pressure and incubation treatments on the consistency of the tomato juice were measured by determining the serum to pellet ratio and the Bostwick value of the treated tomato juice. The serum to pellet ratio and Bostwick value were also determined for non-treated tomato juice, as a control.

25

30 Serum to pellet ratio is a measure of the volume occupancy of the insoluble solids component of tomato juice and was determined by centrifuging the juice (3000g, 10 minutes) and then weighing the resultant serum (solubles) and pellet (insolubles) fractions to determine the ratio. The higher the level of insolubles (ie the lower the serum to pellet ratio), the thicker the sample and the higher its consistency.

35

Bostwick is the standard industry measurement for tomato product viscosity. To obtain juice Bostwick values, the serum was removed from the pellet by centrifugation as above and then added back to 70% of the original sample weight. The measurements were made in a standard Bostwick viscometer. The sample was placed in the

levelled viscometer, the shutter opened and a timing clock started. The distance (in centimetres) at 20°C over which the sample flowed in 30s was recorded. The lower the Bostwick value, the thicker the sample and the higher its consistency.

The serum to pellet ratios and the Bostwick values determined are shown in Table 3.

Table 3

Treatment	Serum:pellet	Bostwick (cm)
No treatment (control)	3.2	8.1
400MPa 10 mins + 40°C 30 mins	2.7	10.1
400MPa 10 mins + 60°C 30 mins	2.8	10.3
No treatment (control)	2.8	8.8
600MPa 20 mins + 40°C 30 mins	1.4	6.1
600MPa 20 mins + 60°C 30 mins	1.2	5.9

From this example and example 1, the following conclusions can be drawn.

Tomato juice given a UHP treatment of 400MPa for 10 minutes still has 100% of its PG activity remaining (Table 1). Even after a thermal incubation of 60°C for 30 mins there is still 45% PG activity remaining (Table 2). During the mild thermal incubation this remaining activity is able to depolymerize pectin in the juice

which is reflected in the higher Bostwick values and serum to pellet ratios (ie lower consistencies) of the treated samples.

5 Tomato juice given a UHP treatment of 600MPa for 20 minutes has no PG activity remaining after treatment (Table 1). However PME is unaffected by this treatment and during the mild thermal incubation PME activity is stimulated (PME has a temperature optimum close to 60°C) to catalyse the removal of methyl ester groups from pectin molecules resulting in the pectin molecules being able to associate with each other via cation cross-links to provide a decrease in Bostwick values and serum to pellet ratios, ie. an increase in consistency.

### 15 Example 3

20 The juice colours of the final samples of example 2 were measured using an Ultrascan XE™; this is a machine for measuring colour available from Hunterlab, USA. The measurements are shown in Table 4 below; the values given are the ratios of the Hunterlab "a" and "b" scores; these scores are used by Hunterlab to describe food colour; "a" is a red to green scale; "b" is a yellow to blue scale.

25 Table 4

Treatment	a/b ratio
No treatment (control)	1.89
400MPa 10 mins + 40°C 30 mins	1.96
400MPa 10 mins + 60°C 30 mins	1.85

600MPa 20 mins + 40°C 30 mins	2.00
600MPa 20 mins + 60°C 30 mins	1.93

5

From these results it can be concluded that UHP treatment in combination with mild thermal incubation has very little effect on tomato colour. This is beneficial as it is desirable to retain as much of the original fresh tomato red colour as possible in the final product. In contrast, the prior art heat processes lead to the generation of brown colours due to the Maillard reaction.

10

#### Example 4

15

A sample of 150g of tomato dice was subjected to a pressure of 500 MPa for 15 mins, incubated in a water bath for 30 mins at 42°C and pasteurised, all in accordance with the method of example 1. Qualitative volatile analysis was then carried out on the sample using a Tekmar™ dynamic headspace (purge & trap) apparatus (model LSC2000 available from Interscience). Four grams of the sample was purged for 10 minutes with helium at 45°C. The volatiles were trapped on Tenax™ absorbing material and cryofocussed before injection into a gas chromatograph. Qualitative information was obtained by means of a dual column system, using chromatography columns of differing polarities (namely models CPSil05 and CPSil13 available from Chrompack), and a reference database. Semi-quantitative information was obtained by comparing peak heights and areas. This analysis was also carried out on a sample of untreated tomato dice pasteurised in tomato juice, as a control.

20

25

30

35

Relative peak heights for selected volatiles in the samples are shown in Table 5. 3-methylbutanal, hexanal,

cis-3-hexenal and 6-methyl-5-hepten-2-one are important for both fresh and processed tomato aroma (Buttery RG, 1993, In Flavour Science: Sensible Principles and Techniques, ACS, 259-286). Cis-3-hexen-1-ol is important for fresh tomato aroma.

Table 5

Volatile	Peak height relative to untreated control	
	untreated (control)	UHP 500MPa 15mins + 42° 30mins
3-methylbutanal	100	135
2-methylbutan-1-ol	100	100
hexanal/cis-3-hexenal	100	350
cis-3-hexen-1-ol	100	125
1-hexanol	100	125
6-methyl-5-hepten-2-one	100	150

This shows that the UHP treatment resulted in an increase in 3-methylbutanal, hexanal, cis-3-hexenal, cis-3-hexen-1-ol, 1-hexanol and 6-methyl-5-hepten-2-one.

It can be concluded from Table 1 that a UHP treatment of 500MPa for 15 minutes is sufficient to inactivate PG, but not sufficient to inactivate peroxidase or PME. Without wishing to be bound by theory, the inventors believe that the presence of peroxidase during the incubation step is responsible for the generation of free radicals which may be involved in the oxidation of lipids and carotenoids,



to generate most of the volatiles listed in Table 5.

5 The results of this experiment show that UHP treatment  
can be used to produce tomato dice with altered aroma  
volatile profiles.

CLAIMS5

1. A process for providing a tomato-based product comprising the following steps:

a) applying ultra high pressure to a tomato piece, such that polygalacturonase is inactivated and pectinmethylesterase is not inactivated;

10

b) incubating the tomato piece with endogenous pectinmethylesterase to achieve a desired consistency; and

15

c) inactivating the pectinmethylesterase.

2. A process as claimed in claim 1, wherein, in step a, peroxidase is not inactivated.

20

3. A process as claimed in claim 2, wherein, in step c, peroxidase is inactivated.

4. A tomato-based product prepared in accordance with the process claimed in any preceding claim.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 97/01265

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A23L1/212 A23L1/06 A23L1/23 A23L3/015

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 11600 A (K.LILJA) 4 May 1995 see page 1, paragraph 2; claims 1,4-6 see page 2, paragraph 2 see page 3, paragraph 3 ---	1-4
A	EP 0 480 422 A (TOPPAN PRINTING) 15 April 1992 see claims ---	1-4
A	WO 94 28745 A (ASEA BROWN BOVERI) 22 December 1994 see claim 1 see page 1, paragraph 3 - page 2, paragraph 1 --- -/-	1-4

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*&\* document member of the same patent family

Date of the actual completion of the international search

4 July 1997

Date of mailing of the international search report

14.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Van Moer, A

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 97/01265

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>"High pressure and biotechnology" 1992, INSERM, FRANCE XP000600625 cited in the application pages 195-209 "Effects of high hydrostatic pressure on food constituents: an overview" J.-C.CHEFTEL see page 199, paragraph 4 see page 202, paragraph 7 - paragraph 8 -----</p>	1-4

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/01265

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9511600 A	04-05-95	SE 500905 C AU 8007094 A SE 9303499 A	26-09-94 22-05-95 26-09-94
EP 480422 A	15-04-92	JP 4148666 A JP 7102119 B CA 2051849 A,C DE 69116070 D DE 69116070 T US 5328703 A	21-05-92 08-11-95 13-04-92 15-02-96 19-09-96 12-07-94
WO 9428745 A	22-12-94	SE 501310 C AU 7012094 A BR 9406804 A EP 0701406 A SE 9302035 A US 5579682 A	16-01-95 03-01-95 19-03-96 20-03-96 12-12-94 03-12-96

THIS PAGE BLANK (USPTO)